


Immunopathogenesis of HIV Infection



During the last 5 years, major advances have been made in our understanding of the pathogenesis of human immunodeficiency virus (HIV) disease and in the development of new potent antiviral agents. With regard to HIV pathogenesis, several recent observations have not only changed our perspectives of HIV disease, but have been critical for the design of therapeutic strategies.

in the immunopathogenesis of HIV infection. A number of theories have been proposed to explain the depletion of CD4+ T cells, ranging from direct killing of Adv Exp Med Biol. 1996;402:165-70. Immunopathogenesis of HIV infection: role of alcohol and HIV peptides. Nair MP(1), Schwartz SA. Author information: Highlights: This concise collection touches on the highlights of immunopathogenesis in HIV infection. Tables and helpful cartoons illustrate theAbstract The rate of progression of HIV disease may be substantially different among HIV-infected individuals. Following infection of the host with any virus, theThis 2-part Medscape clinical update, Immunopathogenesis and Immune Response in HIV Infection (part 1) and Immune Reconstitution and ImmunotherapyImmunopathogenesis of HIV infection. Author information: (1)National AIDS Research Institute, G-73, MIDC, Bhosari, Postbox 1895, Pune 411 026, India. However, adaptive immune response is the most critical component of immune system for control of HIV infection.The progressive and irreversible destruction of the immune system represents the hallmark of HIV infection. Even though this process is directly related to theThe Immunopathogenesis of HIV Infection. ZEDA F. ROSENBERG AND ANTHONY S. FAUCI. National Institute of Allergy and Infectious Diseases.,. NationalSummary:HIV may cause CD4-positive T cell depletion not only by direct infection but by causing damage to normal maintenance mechanisms. Findings inImmunopathogenesis of HIV infection. Author information: During primary HIV infection, a burst of viremia occurs that disseminates virus to the lymphoid organs. A potent immune response ensues that substantially, but usually not completely, curtails virus replication.In vivo the reservoir for HIV infection in the peripheral blood is the CD4+ T cell, whereas in other tissues the monocyte/macrophage may play a substantial role. As disease progresses in HIV-infected individuals, the viral burden in the peripheral blood CD4+ T cells increases.Immunopathogenesis of HIV Infection. transparent Pathogenesis of Immune Deficiency in HIV Infection Predictors of Immune Deterioration in HIV InfectionThe human immunodeficiency virus (HIV) has a unique capacity to infect cells of the human immune system, leading eventually to frank immunodeficiency.Enferm Infecc Microbiol Clin. 2011 Mar29(3):216-26. doi: 10.1016/.2011.01.006. [Immunopathogenesis of HIV infection]. [Article in Spanish]. Alcami J(1)in the immunopathogenesis of HIV infection. A number of theories have been proposed to explain the depletion of CD4+ T cells, ranging from direct killing of On Aug 30, 2012, Chiara Cerini (and others) published the chapter: Immunopathogenesis of HIV infection in the book: Principles and PracticeDuring the last 5 years, major advances have been made in our understanding of the pathogenesis of human immunodeficiency virus (HIV) disease and in theAnnu Rev Immunol. 1995;13:487-512. New concepts in the immunopathogenesis of HIV infection. Pantaleo G(1),

Fauci AS. Author information: (1)Laboratory of